




The Chemical Company



For weed control in winter dormant Bermudagrass pasture and hay fields

This supplemental label expires on December 31, 2011.

EPA Reg. No. 241-418

Active Ingredient*:

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine 38.7%

Other Ingredients: 61.3%

Total: 100.0%

*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

OBSERVE AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS, PRECAUTIONARY STATEMENTS, MIXING AND APPLICATION INSTRUCTIONS, AND FOLLOWCROP RESTRICTIONS ON THE Prowl® H₂O herbicide MAIN CONTAINER LABEL BEFORE USING.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. When following the instructions on this supplemental label, the user must have this supplemental label and the entire **Prowl H₂O**, EPA Reg. No. 241-418, container label in possession at the time of pesticide application.

Prowl H₂O is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Uniformly apply **Prowl H₂O** with properly calibrated ground equipment in 10 or more gallons of water or 20 or more gallons of liquid fertilizer per acre, or in 5 or more gallons per acre of water by air. Use sprayers with constant agitation and equipped with nozzles that provide accurate and uniform application.

Adequate rainfall or overhead irrigation after application prior to weed seedling germination will provide the most effective weed control.

Endangered Species Protection

If endangered species occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave untreated buffer zone of 200 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE fine to medium/coarse nozzles.
- If applied by air, leave an untreated buffer zone of 170 feet. Must use straight stream nozzles (D-6 or larger); wind can be no more than 8 mph, and release height must be 15 feet or less.

To determine whether your county has an endangered species, consult the website <http://www.epa.gov/espp/usa-map.htm>.

Endangered Species Bulletins may also be obtained from extension offices or state pesticide agencies. If the bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered species occur in the area to be treated.

SPRAY DRIFT

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interaction of many equipment-related and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. It is the responsibility of the applicator to avoid spray drift onto nontarget areas.

Application Instructions

Prowl® H₂O herbicide may be applied by ground or chemigation, or by air.

Prowl H₂O may be applied in fields of Bermudagrass grown for hay production and/or grown in pastures for livestock grazing. **Apply Prowl H₂O only to established Bermudagrass** (defined as planted in the fall or spring which has gone through a first cutting/mowing) **when in winter dormancy**.

Apply **Prowl H₂O** prior to target weed germination. Uniformly apply at a broadcast rate of 1.1 to 4.2 quarts of **Prowl H₂O** per acre in a single application. **Prowl H₂O** may also be applied in two split applications, with 1/2 the seasonal application rate applied at the onset of winter dormancy followed by the other 1/2 the seasonal application rate applied just prior to spring greenup. **DO NOT** exceed a cumulative total of 4.2 quarts of **Prowl H₂O** per acre in any one crop season.

Use the higher application rate of **Prowl H₂O** where more dense infestations of targeted annual grasses and annual broadleaf weeds are anticipated or when a longer duration of residual weed control is desired.

Prowl H₂O may be applied in a sequential use program or as a tank mix with other registered herbicides that control emerged weeds.

Prowl H₂O may cause temporary injury to Bermudagrass stands. Disease, extremely cold weather, drought, extensive frost heaving, low or high pH, or salinity may weaken stands and make the crop more susceptible to herbicidal injury.

Chemigation Applications

Prowl H₂O may be applied through sprinkler irrigation systems. Follow all recommendations, special instructions and precautions in the general section covering **Chemigation** in **Spraying Instructions** of the **Prowl H₂O** container label. Uniform distribution of **Prowl H₂O**-treated irrigation water is the sole responsibility of the applicator and is required to avoid crop injury, lack of herbicide effectiveness, or illegal pesticide residues in the crop.

Tank Mixes

Prowl H₂O may be tank mixed with other herbicides labeled for use in Bermudagrass fields. BASF recommends testing **Prowl H₂O** tank mixes on a small portion of the target crop to determine if damage is likely to occur.

Physical incompatibility, reduced weed control, or crop injury may result from mixing Prowl H₂O with other pesticides, additives, or fertilizers.

Applications of postemergence herbicides may cause crop injury, such as stunting or chlorosis of Bermudagrass. Consult with your local BASF dealer regarding local tank mix options. Always perform a mixing test to check the compatibility of **Prowl H₂O** with all potential tank mix partners and fertilizers. Follow all precautions and restrictions on the labels of all products applied in combination with **Prowl H₂O**. Always follow the most restrictive label.

Restrictions and Limitations

- **DO NOT** apply **Prowl H₂O** if surface water is present in the field.
- **DO NOT** exceed a cumulative total of 4.2 quarts of **Prowl H₂O** per acre per year.
- **DO NOT** harvest Bermudagrass hay until **60 days** after treatment with **Prowl H₂O**.
- **DO NOT** harvest for forage or allow livestock to graze Bermudagrass until **45 days** after treatment with **Prowl H₂O**.
- Use only on Bermudagrass grazing areas that are controlled/fenced and livestock are excluded for a minimum restriction period of 45 days after treatment with **Prowl H₂O**.
- Use of **Prowl H₂O** on rangeland is prohibited.
- See the main label for directions for Rotational Crops. For crops which can be rotated the following year after application of **Prowl H₂O**, the minimum plant back interval is 270 days after treatment. Follow the most restrictive language for Rotational Crops in this supplemental label and the main product label.

Weeds Controlled

Prowl H₂O applied according to this label, prior to weed germination, will control sandbur in addition to the weeds listed on the **Prowl H₂O** container label (see **Table 1. Weeds Controlled**). If weeds have germinated prior to the application of **Prowl H₂O**, it may be necessary to apply a herbicide labeled for postemergence activity on those weeds. Weed control may be enhanced by using sequential applications of other registered herbicides.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF. 1108

Prowl is a registered trademark of BASF.

© 2010 BASF Corporation.
All rights reserved.

000241-00418.20090707b.NVA 2009-04-195-0129

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


The Chemical Company